

Table 24. Years of Potential Life Lost Due to All Causes and Selected Cancers by Sex and by Race, Florida, 2009

| | Florida (1) | | Female | | Male | | Black | | White | |
|---------------|-------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| | Years | Percent | Years | Percent | Years | Percent | Years | Percent | Years | Percent |
| All Causes o | 1,311,870 | -- | 490,709 | -- | 820,845 | -- | 284,733 | -- | 994,650 | -- |
| All Cancers | 293,053 | 100 | 136,515 | 100 | 156,464 | 100 | 45,516 | 100 | 241,104 | 100 |
| Childhood C | 4,581 | 1.6 | 2,108 | 1.5 | 2,473 | 1.6 | 1,172 | 2.6 | 3,081 | 1.3 |
| Lung & Bron | 75,103 | 25.6 | 31,365 | 23.0 | 43,693 | 27.9 | 8,117 | 17.8 | 65,934 | 27.3 |
| Prostate | 4,919 | 1.7 | | | 4,945 | 3.2 | 1,225 | 2.7 | 3,617 | 1.5 |
| Breast | 27,976 | 9.5 | 27,711 | 20.3 | | | 6,028 | 13.2 | 21,155 | 8.8 |
| Colorectal | 25,590 | 8.7 | 10,707 | 7.8 | 14,882 | 9.5 | 4,776 | 10.5 | 20,173 | 8.4 |
| Bladder | 4,115 | 1.4 | 957 | 0.7 | 3,158 | 2.0 | 386 | 0.8 | 3,684 | 1.5 |
| Head & Necl | 8,733 | 3.0 | 1,841 | 1.3 | 6,892 | 4.4 | 1,342 | 2.9 | 7,354 | 3.1 |
| Non-Hodgkir | 9,825 | 3.4 | 3,450 | 2.5 | 6,375 | 4.1 | 1,921 | 4.2 | 7,614 | 3.2 |
| Melanoma | 6,670 | 2.3 | 2,182 | 1.6 | 4,488 | 2.9 | | | 6,579 | 2.7 |
| Ovary | 7,504 | 2.6 | 7,504 | 5.5 | | | 680 | 1.5 | 6,438 | 2.7 |
| Cervix | 5,226 | 1.8 | 5,226 | 3.8 | | | 1,111 | 2.4 | 4,027 | 1.7 |
| All Other Cai | 117,366 | 40.0 | 45,573 | 33.4 | 71,766 | 45.9 | 19,863 | 43.6 | 94,529 | 39.2 |

(1) Florida and All Cancer totals include years lost in persons of "Other" and unknown races, and unknown sex, males with breast cancer, and blacks with melanoma.

Source of data: Office of Vital Statistics

(2) Years lost to childhood cancers are included in totals for specific cancer sites.